

## Real estimates of mortality following COVID-19 infection

As of March 1, 2020, 79 968 patients in China and 7169 outside of China had tested positive for coronavirus disease 2019 (COVID-19).<sup>1</sup> Among Chinese patients, 2873 deaths had occurred, equivalent to a mortality rate of 3.6% (95% CI 3.5–3.7), while 104 deaths from COVID-19 had been reported outside of China (1.5% [1.2–1.7]). However, these mortality rate estimates are based on the number of deaths relative to the number of confirmed cases of infection, which is not representative of the actual death rate; patients who die on any given day were infected much earlier, and thus the denominator of the mortality rate should be the total number of patients infected at the same time as those who died. Notably, the full denominator remains unknown because asymptomatic cases or patients with very mild symptoms might not be tested and will not be identified. Such cases therefore cannot be included in the estimation of actual mortality rates, since actual estimates pertain to clinically apparent COVID-19 cases.

The maximum incubation period is assumed to be up to 14 days,<sup>2</sup> whereas the median time from onset of symptoms to intensive care unit (ICU) admission is around 10 days.<sup>3,4</sup> Recently, WHO reported that the time between symptom onset and death ranged from about 2 weeks to 8 weeks.<sup>5</sup>

We re-estimated mortality rates by dividing the number of deaths on a given day by the number of patients with confirmed COVID-19 infection 14 days before. On this basis, using WHO data on the cumulative number of deaths to March 1, 2020, mortality rates would be 5.6% (95% CI 5.4–5.8) for China and 15.2% (12.5–17.9) outside of China. Global mortality rates over time using a 14-day delay estimate are shown in the figure, with a curve that levels off to a rate of 5.7% (5.5–5.9), converging with the current WHO estimates. Estimates will increase if a longer delay between onset of illness and death is considered. A recent time-delay adjusted estimation indicates that mortality rate of COVID-19 could be as high as 20% in Wuhan, the epicentre of the outbreak.<sup>6</sup> These findings show that the current figures might underestimate the potential threat of COVID-19 in symptomatic patients.

We declare no competing interests.

\*David Baud, Xiaolong Qi, Karin Nielsen-Saines, Didier Musso, Léo Pomar, Guillaume Favre  
david.baud@chuv.ch

Materno-fetal and Obstetrics Research Unit, Department Woman-Mother-Child, Lausanne University Hospital, 1011 Lausanne, Switzerland (DB, LP, GF); CHESS Center, The First Hospital of Lanzhou University, Lanzhou, Gansu, China (XQ); Division of Pediatric Infectious Diseases, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA (KN-S); Aix Marseille Université, Institut de Recherche pour le Développement, Assistance Publique-Hôpitaux de Marseille, Service de Santé des Armées, Vecteurs—Infections Tropicales et Méditerranéennes (VITROME), Institut Hospitalo-Universitaire—Méditerranée Infection, Marseille, France (DM); and Laboratoire Eurofins—Labazur Guyane, French Guiana, France (DM)

- 1 WHO. Coronavirus disease 2019 (COVID-19). Situation Report—41. March 1, 2020. [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200301-sitrep-41-covid-19.pdf?sfvrsn=6768306d\\_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200301-sitrep-41-covid-19.pdf?sfvrsn=6768306d_2) (accessed March 5, 2020).
- 2 Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20–28 January 2020. *Euro Surveill* 2020; **25**: 2000062.
- 3 Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; **395**: 497–506.
- 4 Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020; published online Feb 7. DOI:10.1001/jama.2020.1585.
- 5 WHO. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19). Feb, 2020. <https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf> (accessed March 5, 2020).
- 6 Mizumoto K, Chowell G. Estimating the risk of 2019 novel coronavirus death during the course of the outbreak in China, 2020. *medRxiv* 2020; published online Feb 23. DOI:10.1101/2020.02.19.20025163 (preprint).




*Lancet Infect Dis* 2020

Published Online

March 12, 2020

[https://doi.org/10.1016/S1473-3099\(20\)30195-X](https://doi.org/10.1016/S1473-3099(20)30195-X)



**Figure:** Global COVID-19 mortality rates (Feb 11 to March 1, 2020)

Current WHO mortality estimates (total deaths divided by total confirmed cases), and mortality rates calculated by dividing the total number of deaths by the total number of confirmed cases 14 days previously.